

---

Женева, 19 ноября - 7 декабря 2001 года

**СПРАВОЧНЫЙ ДОКУМЕНТ О НОВЫХ НАУЧНО-ТЕХНИЧЕСКИХ ДОСТИЖЕНИЯХ,  
ИМЕЮЩИХ ОТНОШЕНИЕ К КОНВЕНЦИИ О ЗАПРЕЩЕНИИ РАЗРАБОТКИ,  
ПРОИЗВОДСТВА И НАКОПЛЕНИЯ ЗАПАСОВ БАКТЕРИОЛОГИЧЕСКОГО  
(БИОЛОГИЧЕСКОГО) И ТОКСИННОГО ОРУЖИЯ И ОБ ИХ УНИЧТОЖЕНИИ**

Подготовлен секретариатом

1. В пункте 22 своего доклада (BWC/CONF.V/PC/1) Подготовительный комитет пятой Конференции государств - участников Конвенции о запрещении разработки, производства и накопления запасов бактериологического (биологического) и токсинного оружия и об их уничтожении по рассмотрению действия Конвенции решил предложить государствам-участникам, желающим сделать это, в том числе правительствам-депозитариям, представить в секретариат информацию о новых научно-технических достижениях, имеющих отношение к Конвенции. Эта информация должна охватывать сферы применения таких достижений, а также отражать их значимость для различных аспектов Конвенции.
2. Настоящий документ содержит информацию, предоставленную государствами-участниками в секретариат по состоянию на 26 октября 2001 года согласно пункту 22 доклада Подготовительного комитета.

Соединенное Королевство Великобритании и Северной Ирландии

**A. Введение**

1. Подготовительный комитет пятой Конференции государств - участников Конвенции о запрещении разработки, производства и накопления запасов бактериологического (биологического) и токсинного оружия и об их уничтожении по рассмотрению действия Конвенции (КБТО) предложил государствам-участникам представить в секретариат информацию о новых научно-технических достижениях, имеющих отношение к Конвенции. Эта информация должна охватывать сферы применения таких достижений, а также отражать их значимость для различных аспектов Конвенции.

2. Аналогичные предложения были сформулированы для предыдущих обзорных конференций, хотя применительно к первым двум такое обращение было адресовано только трем депозитариям. Это приглашение было рассчитано на информационное обеспечение обзорной Конференции в качестве подспорья в связи с ее задачей в соответствии со статьей XII Конвенции:

*"чтобы иметь уверенность, что цели, изложенные в преамбуле, и положения Конвенции... осуществляются. При таком рассмотрении должны быть приняты во внимание все новые научно-технические достижения, имеющие отношение к Конвенции".*

3. Соединенное Королевство представляло такие справки всем предыдущим обзорным конференциям. В нашей справке, представленной в 1996 году четвертой обзорной Конференции, были рассмотрены достижения в научной, медицинской, сельскохозяйственной сферах с предыдущей (1991 года) обзорной Конференции. В ней также обсуждены последствия для сферы охвата Конвенции и для баланса между потенциалом злоупотреблением наукой и техникой, с одной стороны, и потенциалом профилактического, защитного или иного мирного применения - с другой. Кроме того, были идентифицированы проблемы, которыми, пожалуй, надо было бы заняться в рамках текущей работы Специальной группы, уполномоченной на составление юридически связывающего документа по укреплению Конвенции. Такие проблемы включали разработки и методы, которые могли бы сказаться на конструировании режима объявлений по будущему протоколу к Конвенции.

4. В справке Соединенного Королевства за 1996 год отмечался рост потенциала к широкомасштабному производству потенциальных биологических оружейных (БО) агентов и к генетической модификации агентов и токсинов с целью повышения их оружейной полезности. Вместе с тем, пожалуй, ничто не говорило о том, чтобы распространитель, располагающий мощностями для избрания курса на создание технически передовых вооружений, мог пойти на это со сколь-либо более высокой степенью вероятности, с учетом возросшего риска неудачи или обнаружения, в частности по будущему протоколу к Конвенции. Было также очевидно, что с прежней обзорной Конференции возрос и потенциал в плане защитных мер.

#### **В. Научно-технологический обзор**

5. В настоящей справке воспроизведен широкий подход предыдущего представления, хотя в ней не предпринимается попыток отметить разработки, которые могли бы сказаться на какой-либо из мер укрепления доверия или контроля за соблюдением, которые могли

бы быть предложены в будущем для Конвенции. Сам же размах достижений в сфере генетики, вероятно, мало кого удивит, учитывая степень просвещенности - а порой и озабоченности - общественности по поводу трансгенных животных и растений и генно-инженерных продуктов. Мы сочли важным вновь разобрать прогресс генетических познаний и прикладного применения в связи с микроорганизмами, токсинами и инфекционными заболеваниями, который с обзорной Конференции 1996 года отличался взрывоподобным развитием с точки зрения достижений и потенциала. Вместе с тем имеется и ряд других бурно развивающихся сфер, таких, как протеомика и биоинформатика, которые также заслуживают того, чтобы быть принятыми во внимание в контексте потенциала - будь то благого или недоброго - в связи с положениями Конвенции.

6. Детали такого обзора излагаются в приложении по следующим рубрикам:

- геномика и протеомика
- биоинформатика
- проект "Геном человека" и человеческое разнообразие
- генотерапия
- вирулентность и патогенность
- вакцины и новаторские терапии
- экспрессия рекомбинантных белков
- токсины и другие биоактивные молекулы
- технологии обнаружения и идентификации
- модели инфекционных заболеваний человека
- уничтожение натуральной оспы
- резистентность к лекарственным препаратам
- заболевания в сельском хозяйстве
- борьба с вредителями в сельском хозяйстве
- глобальные инициативы по борьбе с заболеваниями
- прикладная молекулярная биология и культуры
- тенденции в сфере технологий продуцирования белка
- международное сотрудничество и биобезопасность: деятельность по Конвенции о биоразнообразии
- способы доставки агентов или токсинов
- использование патогенов для контроля сорняков и "криминальных" культур
- биоремедиация: уничтожение материала
- противодействие угрозе биотерроризма
- воздействие вступления в силу КХО.

### С. Анализ и рекомендации

7. Приводимый в приложении обзор научно-технологических разработок с четвертой обзорной Конференции 1996 года свидетельствует о том, что в ряде областей достигнут значительный прогресс, который дает представление о генетической, структурной и функциональной основе микроорганизмов и токсинов, а также о количестве и изощренности методов, позволяющих производить целенаправленные изменения свойств микроорганизмов и токсинов. Сдвиги в сфере генетики нередко характеризуются как взрывоподобные, но в этот период бурным развитием были отмечены и другие области. А кое-какие из них даже получили новые этикетки: появились, например, такие новые области, как протеомика и биоинформатика. В биотехнологических секторах многих стран стало обыденным делом продуцирование молекул, встраиваемых генно-инженерным способом в микробные или эукариотные клетки, или же в "трансгенные" растения или животных. Как ожидается, такие сдвиги будут ускоряться, и в следующий период мы станем свидетелями расширения представлений о функциональных взаимоотношениях внутри микробной клетки.

8. С последней обзорной Конференции также заметно возросли возможности злоупотребления токсинами и молекулами пептидных биорегуляторов. Уже имеются в наличии сведения о структуре и функции токсинов и их субъединиц, а сдвиги в рекомбинантной инженерии и системах экспрессии позволили произвести экспрессию ряда токсинов в гетерологичных хозяйских клетках. Таким образом можно легко производить массовое продуцирование и большого числа других токсинов. Поступают сообщения о рекуперации пептидов из тонновых количеств трансгенных растений. Тем самым сдвиги в сфере биотехнологии создают потенциал для злоупотребления пептидными биорегуляторами после продуцирования в рекомбинантных системах. Сдвиги в векторной доставке открывают перспективы использования вирусных или бактериальных векторов для прямой доставки токсина/биорегулятора к мишени либо с целью генетической передачи хозяину, либо с целью прямого продуцирования вектором. Сдвиги в рекомбинантном производстве также означают рост потенциала для того, чтобы принимать в расчет возможность оружейного использования прионов. Как полагает СК, обзорная Конференция должна учитывать эти факторы при разборе статьи I.

9. Глобальное распространение знаний в сферах, имеющих отношение к микробиологии, да еще и в сочетании с таким подспорьем, как новые информационные маршруты типа Интернет, означает, что возрастает и число стран, которые располагают учеными, обладающими знаниями и квалификацией для разработки биологического оружия на основе классических потенциальных агентов и токсинов. Но, пожалуй, еще большую озабоченность вызывает то обстоятельство, что распространение знаний и

прикладной практики в таких областях, как геномика, означает, что уже сейчас во все более широком круге стран могли бы быть предприняты попытки произвести усовершенствование биологических оружейных агентов за счет их генетического изменения с целью усиления таких свойств, как жизнестойкость, резистентность к антибиотикам и способность преодолевать специфические методы обнаружения и профилактики. Пока еще слабо уяснены многие аспекты, имеющие отношение к свойствам микроорганизмов, например в рамках взаимоотношения между их различными структурно-функциональными свойствами и природы взаимодействий с хозяином в ходе инфицирования. Однако развитие и распространение познаний открывают новые возможности с точки зрения распространителя БО, которые следует принимать в расчет. По мере роста научных познаний будет возрастать и глобальный потенциал для технически передовых биологических оружейных программ. Судя по уже появившимся данным о человеческом геноме, пока еще не выявлено отличий, которые могли бы быть использованы в качестве основы для "генетического оружия", нацеленного на конкретные этнические группы.

10. Сдвиги в областях, имеющих отношение к Конвенции, вызывают еще две озабоченности. Во-первых, глобальное распространение знаний в сферах науки и технологии повышает потенциал для того, чтобы негосударственные субъекты во все более широком круге стран попытались предпринять производство и распространение патогенов или токсинов в качестве оружия, хотя на практике их успех может ограничить или предотвратить связанный с этим довольно высокий уровень квалификации. В ответ на это несколько правительств приступили к реализации специфических контрмер, включая регламентационные меры с целью предотвратить несанкционированное приобретение патогенов и токсинов в законных коллекциях культур, к созданию инфраструктур в целях преодоления последствий любого применения БО против их населения, а также к закупкам средств обнаружения и вакцинных препаратов. Во-вторых, риск неожиданных результатов работы с генетически модифицированными микроорганизмами не может не возрастать по мере того, как и в развитых, и в развивающихся странах будет возрастать число лабораторий, занимающихся рутинным применением рекомбинантных технологий к микроорганизмам. В качестве предупреждения о рисках непредвиденных последствий, которые могли бы носить катастрофического характера, например в том случае, если бы такие организмы ускользнули из лаборатории, послужил преданный гласности эксперимент с вирусным рекомбинантом мышьяковой оспы, который неожиданно преодолел иммунную защиту хозяйской мыши. А это подчеркивает важность тщательной оценки рисков и надлежащих процедурных и физических мер изоляции. Экологическое высвобождение опасного рекомбинантного микроорганизма могли бы быть сопряжено с возникновением озабоченностей, хотя оно и было бы беспочвенным в контексте КБТО.

11. Для применения биологических агентов и токсинов могут быть приспособлены многие из военных систем доставки, которые, как сообщается, становятся доступными для все более широкого круга стран - как развитых, так и развивающихся. Как ожидается, повышению шансов на эффективную дальнюю доставку биологических оружейных агентов будут по-прежнему способствовать усовершенствования в плане ракетной дальности, точности и конструкции суббоеприпасов.

12. В ряде стран по-прежнему возрастает потенциал и наличие оборонительных мер против БО, причем как с точки зрения полевых систем для обнаружения и идентификации биологических агентов и токсинов, так и с точки зрения профилактических мер, главным образом вакцин. Можно также ожидать, что повышению национальной готовности к биологическим нападениям со стороны других стран будут способствовать национальные программы по борьбе с терроризмом, предусматривающие значительные ассигнования на закупки средств обнаружения и медицинского противодействия. Вместе с тем значительный хронологический разрыв между разработкой практических средств противодействия, таких, как датчики и вакцины, контрастирует с той стремительностью, с какой могут изменяться свойства патогенов и токсинов во многих лабораториях всего мира, которые сейчас могли бы, приступая к нелегальной программе, предпринимать такие модификации. И хотя важно, что в ряде стран происходит совершенствование биологических оборонительных потенциалов, на которых благотворно сказываются технические достижения, рост технической изошенности и сопутствующие расходы могут еще больше затруднить развивающимся странам приобретение эффективных оборонительных средств.

13. Произошло усугубление глобальной пандемии СПИДа, и возрастают опасения по поводу распространения СПИДа в странах, где уже эндемичен ТБ. Рост глобальной мобильности и торговли по-прежнему способствовал неожиданным естественным вспышкам инфекционных заболеваний. Причем такие вспышки могут выходить за рамки нормального идентификационного и терапевтического потенциала страны, ну а что касается, в частности, заболеваний растений и животных, то, даже не отличаясь очень уж высокой заболеваемостью или смертностью, они могут быть сопряжены с колоссальными экономическими издержками. В качестве примера можно сослаться на прямые издержки эпидемии ящура в СК в 2001 году, а также на ее еще более высокие косвенные издержки для такого рода деятельности, как индустрия туризма.

14. Что касается положительной стороны дела, то международные организации, занимающиеся профилактикой заболеваний человека, животных и растений, все больше прибегают к совместной деятельности и поощряют координацию национальных и

региональных программ. Настоящая справка приводит примеры вкладов СК в проекты, нацеленные на борьбу с заболеваниями в развивающихся странах. Согласован Картахенский протокол к Конвенции о биологическом разнообразии; он обеспечит принятие мер, рассчитанных на снижение рисков для биоразнообразия, которые могли бы происходить из трансграничного движения живых (генетически) модифицированных организмов. Новые представления о взаимосвязях между вспышками заболеваний как в разных странах, так и в хронологическом разрезе удалось получить благодаря техническим достижениям в диагностике заболеваний растений и животных, таким, как генетическая типизация. Такие новые диагностические возможности облегчают проведение разграничения между естественными и неестественными вспышками.

15. Продолжаются исследования мирных способов использования микроорганизмов с целью уничтожения загрязнителей в окружающей среде, но эта методика вызывает озабоченности по поводу потенциала переносов нежелательных генов в природные организмы. Все еще ведется рассмотрение одного из способов прикладного использования биоремедиации с целью нейтрализации остатков химических оружейных агентов (ХО) в экологических пробах, хотя этот метод не был воспринят каким-либо из государств - участников КХО для крупномасштабных программ уничтожения химических оружейных агентов. Биоремедиационные технологии явно обладают потенциалом для развития в качестве средства ведения войны или для враждебного применения против материала, имеющего кардинальное значение для нормальной гражданской жизни или для военных операций, такого, как масла, резины и пластики.

16. С обзорного процесса 1996 года расширялось использование микроорганизмов в целях борьбы с сельскохозяйственными вредителями, причем уделяется большое внимание повышению контроля качества при изготовлении препаратов. Продолжались исследования потенциала микробных патогенов с точки зрения борьбы с сорняками, но тут пока еще имеется мало коммерческих способов применения. Ряд объединений подвергают критике программы ООН по разработке патогенов, пригодных для применения в качестве специфических гербицидов с целью уничтожения культур коки, марихуаны и опия, выращиваемых с целью производства незаконных наркотических средств. Подняты, в частности, следующие научные проблемы: непредвиденные генетические переносы и инфицирование растений, помимо намеченных целевых видов. Возникают и вопросы о том, непременно ли было бы приемлемым по положениям статьи I КБТО истребление таких культур с использованием биологических агентов, с учетом того, что сценарии истребления таких культур фигурируют в настоящее время в странах, где отмечаются гражданские беспорядки, а то и внутренние вооруженные конфликты. Как полагает СК обзорной Конференции надлежит обсудить эту проблему с целью установить возможность внесения ясности на этот счет.

17. Значительную диверсификацию претерпело промышленное использование микроорганизмов или их экстрагированных ферментов с целью катализа единичных ступеней в процессах химического изготовления - так называемая "биотрансформация". Можно ожидать, что объекты, использующие такие процессы, будут фигурировать в объявлениях производственных площадок по КХО. Носит и, по-видимому, сохранит ограниченный характер транспарентность, присущая объявлению и рутинным инспекционным процедурам КХО применительно к работам по производству токсинов и других химикатов биологического происхождения. По мнению СК, всякая транспарентность, предусматриваемая будущими мерами по КБТО, должна охватывать гораздо более широкий диапазон токсинов и их аналогов и субъединиц.

18. В ходе различных исследований и консультаций, проведенных СК в качестве информационного подспорья для настоящего обзора, стало ясно, что динамика перемен в сферах науки и техники, имеющих отношение к КБТО, носит гораздо более широкий размах, чем в предыдущий пятилетний период, т.е. в период между третьей и четвертой обзорными конференциями. Ряд сдвигов в сфере научных познаний и практики их применения могли бы быть сопряжены с последствиями для положений КБТО. С учетом быстрых темпов развития науки и техники, СК задается вопросом, осмотрительно ли сохранять пятилетний разрыв между такими оценками по КБТО. СК предлагает предстоящей обзорной Конференции подумать об учреждении механизма для совместной работы государств-участников на более регулярной основе с целью проведения такого рода научно-технических обзоров и с целью рассмотрения любых последствий на необходимом уровне квалификации.



Annex<sup>1</sup>

DETAILED SCIENCE AND TECHNOLOGY REVIEW

Genomics and proteomics

1. The advent of rapid methods for sequencing entire genomes has led to a massive increase in genomic data in recent years. Although sequencing technology is not a novel technique, the last five years have seen the development of faster throughput methods, and this technology is still developing rapidly. As a result, the entire genomes of many viruses and bacteria have now been sequenced. Many of these agents are pathogens and potential biological weapons (BW) agents.
2. To date, 38 bacterial genomes have been published as sequenced, assembled and annotated, and 158 are at various stages of sequencing. In 2001 the genome of *Yersinia pestis*, a potential BW agent that causes plague in humans, was sequenced in the UK. This genome provides evidence for the evolution of the current organism from an enteropathogen ancestor. Genomic information has a wide variety of applications and can underpin studies to develop improved vaccines and antibacterial strategies and to identify antigens as targets for detectors. However, information from genomics that furthers understanding of the mechanisms of microbial pathogenicity also could suggest genetic targets for modulation that could lead to micro-organisms with increased pathogenicity. Genomic information could also inform strategies to defeat vaccines or host immune defences.
3. Although genome sequence data provides information on the entire set of genes a micro-organism possesses, and genomic analysis provides information on the possible function of genes, they do not indicate which of these genes are expressed to produce proteins at any particular time in the life of the organism. Additionally, proteins expressed in organisms are typically modified, for example by phosphorylation or glycolysation, but little can be deduced about this by simply looking at an organism's genome map. Because of the functional complexities of the expression of the genetic potential of a cell, organisms may well have well over an order of magnitude more proteins than genes. The recent finding that the human genome apparently only contains 30,000 - 35,000 genes suggests that much of the complexity in humans lies in the various ways in which proteins are formed and modified.
4. A new field of studies, called proteomics, has evolved as a result of the need to address such issues. Proteomics is the study of the entire set of proteins expressed by the gene sequence

---

<sup>1</sup> Воспроизводится только на английском языке.

of an organism. This allows the study of the total protein complement of a micro-organism, whereas traditionally individual proteins have been studied one by one. In addition, proteomics can allow the characterisation of proteins and of their level of expression under different conditions, and can help elucidate their functional roles. Recent developments have focused on speeding up the procedure of identifying proteins from two-dimensional gels through automation of instrumentation and software. Mass spectrometry has become the technique of choice to identify proteins, and technology developments here have led to lower limits of detection for cellular proteins and increased capabilities for protein and peptide structural studies.

5. Information from proteomic studies could, for example, not only determine which proteins are expressed during microbial infection or disease, but also how proteins interact with each other. Identifying which proteins a target protein interacts with can help in understanding its function. Proteomics can also identify protein markers of disease: this could help in the development of new methods for early diagnosis and detection of diseases such as cancer, or the development of novel therapeutics including vaccines. Possibilities for improving the stability of pathogen or proteins may be exploited in environmental release applications such as bioremediation and the use of biological insecticides and herbicides. Like genomics, such a powerful tool could also be open to misuse for non-peaceful purposes.

6. An example of a technology that is allowing a great increase in analytical capabilities is the use of nucleotide and protein microarrays. Transcriptional profiling using DNA or short oligo-nucleotides arrayed at very high density on chips is an enormously powerful technique which can be used to measure gene expression. These "gene chips" can show which individual genes are switched "on" or "off" under a different conditions. Using such technology the whole genome of a micro-organism can be represented on one chip. Microarrays can be used to ascertain which genes are involved in host invasion by an intracellular pathogen or which genes are expressed when bacteria encounter a host's defence systems. This technology can also provide insights into diseases such as cancer and the genetic basis of other diseases. Protein arrays are receiving widespread attention as analytical tools and can also be used to identify disease biomarkers. Protein chips can be used to study protein-protein interactions and interactions between proteins and drugs. In drug screening, protein arrays can be used to determine which compounds react with target proteins and which react with non-target proteins i.e. those which could potentially cause side effects.

#### Bioinformatics

7. Genome and proteome programmes generate vast amounts of information. The discipline of bioinformatics has emerged and has grown rapidly because of the need to manage this type of information explosion. Bioinformatics is the computer assisted science of processing and dissemination of biological information and involves the collection, storage, management,

classification, integration and retrieval of information in accessible databases. For example, bioinformatics can search for genes of interest making it possible to sort through a genome sequence, select a gene, determine whether it has a known function by comparing it to gene databases, and predict its cellular location and molecular weight.

8. The Internet has made the data and tools for bioinformatics available widely and at low cost, and there are a number of powerful analytical programmes to allow the data for the millions of bases generated during genome sequencing to be mined relatively easily for genes of interest. Software for genome annotation, gene finding, sequence similarity searches, sequence alignments and other analytical functions is constantly being improved. Genetic and protein databases have been growing exponentially over the last few years and protein structural predictions from sequences have become increasingly possible and more accurate as more entries are posted in such databases. Bioinformatics can aid the design of novel drugs and therapeutics, but could equally aid the design of novel toxins or improvements in pathogenicity that could be misused. In the future, bioinformatics linked with high throughput methods for proteome and genome analysis, such as microarrays, will increasingly allow the rapid targeting of biological macromolecules for any purpose, peaceful or otherwise.

#### Human Genome Project and Human Diversity

9. A working draft of the sequence of the human genome, providing greater than 90% coverage of the genome, was published under the Human Genome Project (HGP) in February 2001. The fact that this was achieved ahead of schedule illustrates the rate of progress in such technologies. This draft still has some short gaps and the full sequence will take a further 2-3 years to complete. So far it has been estimated that 30 000 - 35 000 human genes make up the human genome. The potential benefits of this project are enormous, providing a wealth of opportunities to improve diagnosis and understanding of disease, developing gene therapies, and rationalising drug design.

10. The human genome consists of 3 billion nucleotides of DNA of which the majority (99.9 %) is now believed to be identical between individuals. It is the remaining 0.1% which makes us individual, and much work has centred on understanding these differences and their role in disease and response to drugs. The field of pharmacogenetics has developed to study how differences in genes affect the way individuals respond to medicines. Pharmacogenetics could lead to the development of tailor-made medicines and styled therapies for the individual making safer and more effective treatments. One key to understanding this genetic variation is the phenomenon of single nucleotide polymorphism. This is the most common type of genetic variation, occurring once every 1000 base pairs. Many of these base pair changes may have no effect, while others may be involved in disease susceptibility and responses to drugs. In 1999 the Wellcome Trust Foundation set up the SNP Consortium, a \$45 million initiative funded by

the Wellcome Trust and 10 pharmaceutical companies as a non-profit making venture. Their aim is to make this type of information available in the public domain without intellectual property restrictions on academic or commercial researchers accessing this information.

11. The HGP has carried out studies which show that very little extra information in the areas of motif scanning, gene identification, positional cloning, single nucleotide polymorphism mapping and gene structure determination will be gained from the finished sequence as compared to the working draft. This working draft is freely available in public databases and none of the information will be patented.

12. Information from human genomic studies has increased our knowledge of receptors for natural ligands. Such information may provide information on novel receptor targets for modulation of the immune system or central nervous systems, for example. In addition, it may aid the design of molecules intended to mimic natural ligands as a means of modulating regulatory processes. The technology could however be misused to develop novel bioregulators as BW agents.

13. At the beginning of 2000, the company Celera announced their intention to sequence the genomes of 5 individuals, 3 women and 2 men. In an attempt to quantify human diversity at the genome level these individuals were chosen as representing the main ethnic groups of the human race. Results of the analysis were published in the open literature in 2001. Notwithstanding the potential of such projects for scientific and medical benefits, a number of geneticists and medical ethics specialists have raised concerns about the potential for misuse of such information. One type of misuse could be use by a BW proliferator to design a genetic weapon able to target population specific genetic traits. The UK's paper to the 1996 Review Conference drew attention to this potential for genetic weapons, but assessed that genetic weapons could well remain a theoretical possibility. New genetic data provide no indication as yet of differences between ethnic groups that would allow targeting by weapons. Nevertheless, we consider it important that one of the Article I reaffirmations at the Fourth Review Conference stated that Article I applies, *inter alia*, to 'any applications resulting from genome studies'.

#### Gene therapy

14. Gene therapy, the transfer of genetic material for therapeutic purposes, is still not a widely used therapeutic technique. The major challenge associated with this technology remains the gene delivery system. However, the last five years have seen a steady increase in the number of publications about the development of replication competent viral vectors. Three virus groups are being investigated for therapeutic use: retroviruses; adenoviruses; and herpes simplex virus. A more effective delivery system reduces the number of recombinant virus particles needed for

effective during therapy. Such viral vectors would replicate in human cells giving improved tissue dissemination and increased expression levels of any detrimental gene carried by the virus vector.

15. Research procedures using gene therapy are being investigated in animal models to modify inappropriate immune responses associated with conditions such as irritable bowel syndrome. The expression of interleukins delivered by these virus vectors as a means to aid an immune response in cancer treatments has progressed as far as clinical trials. Adenovirus vectors have been used to deliver interleukin 4, a negative feedback chemical messenger of the host immune system.

16. Over the last five years there has been a steady maturation of related technologies such as production and purification methods for viral vectors, new non-viral delivery systems, targeted delivery of therapeutic genes, regulated expression and lung gene therapy techniques. In the long term such technologies may improve sufficiently to be used by a BW aggressor to deliver a foreign gene into human cells.

17. Among the latest technologies, antisense therapy, the use of short spans of nucleic acid to disrupt the expression of disease related gene sequences, is receiving attention as a novel therapy for diseases such as AIDS and Crohn's Disease. Normally, the code of a gene (its sense strand) is transcribed into mRNA which is then translated into a protein. This gene expression can be interrupted either at the stage of transcription or translation by the delivery of an antisense strand of nucleic acid, which by hybridising to either the DNA or mRNA would block the expression of genes and protein products, for example those involved in disease. A related area of interest which is receiving attention is the use of ribozymes as a therapy to alter gene expression. Ribozymes are RNA molecules which can cleave RNA at specific sites. By combining specific antisense targeting with enzymic cleavage of RNA by ribozymes, ribozymes could be used to inhibit protein products associated with disease or those associated with the replication of pathogens in the host.

#### Virulence and pathogenicity

18. Genome sequencing of pathogens has had a significant impact on the understanding of pathogenicity. Pathogenicity islands, where clusters of virulence-associated genes are found, have been shown to be as important as plasmid acquisition in the emergence of pathogens from non-pathogenic ancestors. Many pathogenic strategies, such as Type III secretion systems, have been shown to be widespread amongst bacteria. Evidence is amassing that the virulence of many bacterial pathogens is a complex, multifactorial process requiring the co-ordinated activity of gene products. This regulation of genes is known to be crucial for the survival and virulence of pathogens *in vivo*. In recent years there have been rapid advances in experimental approaches

for the study of bacterial virulence in the host which greatly simplify *in vitro* analysis. High-throughput, global approaches such as signature-tagged mutagenesis and *in vitro* expression technology have been developed. Combined with modern approaches to analysis of gene expression involving DNA micro-arrays, the effect of altered environmental conditions or a mutation on every gene in the bacterium can now be examined.

19. Many virulence studies require an *in vivo* screen in an animal model. Over the last five years there has been an ethical drive in some states to reduce experimental animal use, which has led to the development of alternative models for example using tissue culture or nematode worms. Many studies, ranging from pathogenicity studies to vaccine development will however continue to rely on small animal tests for the immediate future.

20. Advances in genomics have increased our understanding of viral pathogenicity and virus replication in the host. The function of individual genes may be assigned by the expression of individual proteins and analysis *in vitro*, production of attenuated strains, or the generation of recombinant viruses containing foreign or manipulated genes. However, because precise genetic manipulation of viruses is only possible using DNA, DNA copies of RNA viruses must be prepared before their genomes can be manipulated. This process has been aided by recent advances which have increased the fidelity of the polymerase chain reaction (PCR), and is now relatively routine for positive (sense strand) genome RNA viruses. The most significant technical advances have been in reverse genetics, the preparation of a DNA copy of a negative strand (antisense) RNA genome, particularly where the genome is segmented as in the case of influenza virus. This technology has been applied to the highly pathogenic filoviruses to study the function of the virion glycoproteins.

21. Experiments involving the production of recombinant viruses expressing foreign genes have generated unexpected and controversial results. In 2001, a research group described an unexpected lethal effect in mice following their immunisation with orthopoxvirus ectromelia (mousepox, ECTV) into which the murine interleukin-4 (IL4) gene had been inserted. The aim of the insertion was to bias the host immune response and so make the mice infertile - an attempt at contraception.

22. The implications of this mousepox data could be far reaching. ECTV is a close relative of smallpox virus and of vaccinia virus, used in the smallpox vaccine. In addition, ECTV has been adopted as a model virus for smallpox, because it causes a severe and fatal generalised disease in some strains of mice. A parallel can be drawn between some observations made with ECTV in mice and smallpox virus in humans. These results, if extrapolated, could mean that similar experiments with smallpox could result in recombinant viruses with enhanced pathogenicity in humans or the ability to overcome vaccination. Other orthopox viruses such as monkeypox

could be changed in the same way. Unfortunately, such ideas could be taken up in a weapons programme. The authors considered carefully whether it was appropriate to publish the information. Ultimately, it was decided to publish so as to warn others about the potential for experiments with recombinant micro-organisms to have unforeseen results. This emphasises the importance of prior risk analysis and appropriate procedural and physical containment measures.

23. Understanding the pathogenicity of a pathogen is key to constructing an effective defence against disease. However, although an aggressor can use a pathogen without a thorough understanding of its pathogenicity, such knowledge may point to avenues which would increase an agent's effectiveness as a weapon.

#### Vaccines and novel therapies

24. The greatest advances in vaccine research and development have arisen from the availability of genome sequences of pathogens and high-throughput screening methods. Vaccine development requires the identification of unique structures and molecules that can be used to generate immunological protection. Using traditional screening techniques this process can be time consuming and involve the trial and error testing of vaccine candidates. Several groups around the world have now shown that it is possible to predict preferred vaccine designs from genome sequences using bioinformatics software and programs. Whole genome libraries can be screened in animal models by use of methods such as signature tagged mutagenesis. Together, these approaches mean that vaccine discovery now has the potential to proceed much faster and with a better chance of leading to acceptable vaccines.

25. Although conventional sub-unit or killed vaccines are still widely used, recombinant sub-unit vaccines are expected to gradually replace them. But for many disease targets there is a continuing debate over whether highly purified protein antigens will be more effective than a mixture of cells and/or components. Current mixtures tend to be better defined than classical mixed antigen mixtures because some of least of the components are often contributed by recombinants. Pertussis (whooping cough) is an example where a consensus strategy in favour of cellular vaccines is evolving towards a better-defined mixture. Improvements in fermentation technology allow more consistent production of classical vaccines, but on-line monitoring and feedback control of antigen expression during production is still not possible.

26. Recombinant vaccines offer advantages not only in cost of production, yield and safety of the production process but also because of the potential for reduced reactogenicity of the vaccine. Several naked DNA vaccines directed against viral or parasite pathogens including HIV and malaria are now in clinical trials. Vaccine vectors have undergone limited trials in man: for example, salmonella expressing tetanus Hc fragment has been evaluated in a phase 1

trial. However, such live vaccine vectors might equally well be misused to deliver toxins or other biologically active molecules to humans in a weapons format.

27. An area also receiving much attention is the development of more effective vaccine delivery systems. Groups in industry and academia are currently developing a range of improved vaccine adjuvants, with the objectives of enhancing the immune response and tailoring it to the type of protection required. Improvements in vaccine efficacy have been achieved by adding fragments of bacterial toxins to a vaccine. For example, the non-toxic B sub-unit of cholera toxin has proved an effective adjuvant for orally or intranasally delivered vaccines.

28. Microencapsulation technologies are also being applied to improve vaccine delivery. There is an increasing awareness of the advantages of this approach to intranasal delivery. Unfortunately, advances in the intranasal delivery of vaccines could also be misused for the design of improved systems for the airborne delivery of agents and toxins as weapons.

29. The last five years have also seen advances in the re-targeting of microbial toxins to alternative sites of action for therapeutic purposes, and this approach is being adopted in searches for novel drugs for treatment of tumours and. Production of heterologous toxins by the substitution of binding domains with those from non-related proteins has been reported. The incorporation of lectin or antibody into liposomes is being examined as method for targeted toxin delivery to specific cell types in novel therapies. The use of phage display for the production and selection of target specific antibodies for use in immunotoxins has also been widely reported.

30. There is increasing interest in the potential of transgenic plants for use in vaccine production, introducing the foreign gene by means of plant viruses. There have been reports of animal trials with plant vaccines for Hepatitis B virus, rabies virus, respiratory syncytial virus and HIV virus, and clinical trials have probably now started. Much of the work has concentrated on the use of edible plants, with two prospective benefits: that purification can then be minimised or avoided; and that the stimulation of mucosal immunity resulting from oral delivery may provide enhanced protection against infections that invade through the mucosa.

31. There have been reports of a number of national development and procurement programmes for vaccines required for 'consequence management' of biological attacks by state or terrorist actors. Announcements by the Joint Vaccine Acquisition Program (JVAP) in the US Department of Defence in 2000 indicated that it is seeking sources for several vaccines as a stockpile to protect US forces against BW. Vaccine requirements now on the JVAP list now include ricin, tularemia, smallpox, botulinum toxins (multivalent, recombinant), Venezuelan equine encephalitis, and the next generation anthrax vaccine. The new generation of smallpox vaccines use the same vaccinia strain as the old but grown in animal cell lines rather than



recovered after scarification of animals. In the UK, the Ministry of Defence announced in March 2001 that it had received new supplies of anthrax vaccine, and planned to resume its programme of voluntary immunisation against anthrax for armed forces personnel deployed to the Persian Gulf. For many of the putative BW agents, the development of vaccines continues to suffer from the lack of surrogate animal markers of efficacy, and phase III clinical trials cannot be mounted because of the absence of human population groups naturally exposed to the disease.

32. There has been marked progress in the development of anti-viral drugs since the last Review Conference. Acyclovir, one of the most successful drugs against herpes viruses, has been approved in an ester form for oral delivery. Cidofovir is currently used for the treatment of cytomegalovirus retinitis in AIDS patients. It has been shown to have activity *in vitro* against smallpox, but it has the disadvantage of high toxicity, which can lead to renal failure, and thus it is unlikely to be a realistic candidate for use in an outbreak of smallpox.

33. The targets for new candidate antiviral drugs include inhibition of a number of different functions: cell entry; viral protease; viral nucleic acid synthesis and processing; regulatory proteins; and viral release. Many of the most promising new drugs have relatively specific activity against viral agents, which generates the need for development of specific virus diagnostic tests. For antibacterial compounds, there has been continuing development of the older class of broad-spectrum antibiotics, with the objective of extending the spectrum of action and improving their efficacy against organisms resistant to the parent compound. Some new antibacterial drugs tend to have narrow spectrum activity: daptomycin, which acts by disrupting gram positive cell membranes, is an example. The first compound in the new class of oxazolidinones, linezolid, is active against gram positive bacteria and has been particularly useful in treating skin infections and pneumonias.

34. A number of other approaches to antimicrobial therapy are being explored, including immunomodulators such as cytokines and non-specific immunostimulators. An example of the latter is ribostine, an RNA -salt preparation that has been reported to stimulate interferon production and to be beneficial in treating herpes virus and other urogenital infections. To date, the only drug with a proven track record is Interferon-alpha, which is used in AIDS therapy. Other interferons are likely to find applications in the future, perhaps for use in conjunction with antibiotics in treating tuberculosis.

35. In the long term, these technical developments must increase the potential for development of improved vaccines for use to protect armed forces and civil populations against BW attack by state or sub-state actors, and for use in related consequence management.

## Recombinant protein expression

36. Development of recombinant DNA techniques has facilitated the integration of “upstream” gene cloning methods with “downstream” protein purification methods. Production of recombinant proteins has developed significantly over the last five years with a vast choice of cloning and host systems now available. Hosts include bacteria, yeast, plants, filamentous fungi, and insect and mammalian cells. The new developments allow the expression of proteins with increased yield and solubility, tight transcriptional control, optimised codon usage and directed protein localisation. Developments in cloning technologies have included availability of high fidelity thermostable polymerase enzymes, pre-dispensed PCR reagents, ligase free cloning systems and cell free expression systems. Such high throughput systems significantly reduce time scales for the initial steps in protein expression and expression screening.

37. Affinity chromatography purification matrices are used for protein purification to produce high yields of specific product. The area of protein purification process development has seen major advances in instrumentation. There are now a range of platforms that can be used for research laboratory based purification and characterisation of bio-molecules through to scale up and optimisation for industrial production. These instruments offer a high level of automation, with pre-programming that minimises run times and allows method scouting to achieve optimum conditions.

38. An example of a new technology with great potential for directed evolution is applied molecular evolution technology (DNA shuffling). DNA shuffling allows the generation of large libraries of chimeric and mutated hybrids of genes, genomes, plasmids and viruses. Target DNA is randomly fragmented and reassembled to create a large number of recombinants that can be screened for novel or desirable properties. The primary application so far has been the rapid development of proteins for a wide variety of industrial applications such as enzymes with altered substrate affinity and stability. DNA shuffling can also be used to generate novel vaccine antigens for protection against pathogens. Such a technology could be used to develop both toxins and pathogens with altered properties.

39. All these developments mean that proteins previously difficult to produce or extract on any scale can now be expressed relatively easily. With this technology it is relatively easy to produce toxins and toxin sub-units rapidly on a large scale using basic laboratory skills. Although many of these recombinant proteins may be used as vaccines and other therapeutics, there is also the potential to produce toxins and other protein agents with enhanced properties for offensive use as weapons, and on scales which are far beyond those allowed by the natural sources of the molecules.

## Toxins and other bioactive molecules

40. The structure of many more toxins has now been elucidated, and the different functions of their sub-units are increasingly known. Most microbial toxins (or their sub-units) have been cloned into micro-organisms and this technology is now well established. Cloning and expression of animal toxins is as yet not as advanced, but the past five years have seen a large increase in studies where animal venoms have been expressed in *Escherichia coli*, *Pichia pastoris* or baculovirus expression systems. The range of expression vectors, host strains, purification systems and protein engineering systems available have greatly increased the potential to express toxins with enhanced toxicity.
41. Research into scorpion, snake and spider venoms is widely reported in the literature with the emphasis either on anti-venom research or use as tools for studying membrane transport systems. The potential use of the animal toxins as BW agents has previously been thought unlikely due to the difficulty of producing recombinant toxins in appreciable quantities. However, this could change as the cloning of more animal toxins continues.
42. Increasing attention is being given to developing medical applications of clostridial neurotoxins. The therapeutic use of botulinum toxin has been extended beyond its initial application in treating squint, to include a variety of neurological disorders associated with involuntary spasmodic contraction of muscles, cosmetic treatment of frowns and wrinkles, and to facilitate wound healing after surgery. Specific therapeutic uses for clostridial neurotoxins are being proposed based either on delivery and membrane binding functions of the heavy chain or on the light chain function of intracellular inhibition of neurosecretion. Botulinum neurotoxin heavy chain offers opportunities for selective macromolecule drug delivery to the peripheral nervous system, and tetanus heavy chain to the central nervous system, which could lead to treatments for chronic pain.
43. The period has brought an enormous increase in knowledge about the structure of cellular receptors in man, animal and plants, and about the molecular basis of bioregulation, including immunomodulation. Many of the small bioactive molecules are proteins or peptides that in principle could be easily produced in recombinant micro-organisms or in transgenic animals or plants, making production on weapon scales increasingly feasible. There is also the possibility of misusing microbial vectors to deliver bioregulator genes or to cause bioregulator production directly in the warfare target.

## Detection and identification technologies

44. Recent developments in molecular techniques particularly in the PCR have improved the potential for the rapid detection and identification of micro-organisms, including in the context of defence against biological attack. PCR techniques now allow highly specific and sensitive assays to be developed as a routine. With more rapid thermal cycling, PCR reaction times have been reduced from 2-3 hours to as little as 10 minutes. Although not yet available commercially, "DNA chips", microfabricated silicon PCR vessels that use the thermal properties of small volumes and silicon, have the potential to increase speeds even further. The second major improvement has been the advent of PCR instruments that combine thermal cyclers with a fluorimeter that can detect amplification by the incorporation of fluorescent components into the PCR chemistry. Taking fluorescent measurements during the amplification and thus measuring the reaction kinetics reduces analysis time and detection thresholds. These systems are now available, including versions for military and law enforcement field applications. Many of the developments improve the automation of PCR. Reagents are also now available in more convenient formats such as pre-prepared reaction cocktails and freeze dried "beads". Much of the commercial development has involved the use of robotics for laboratories needing to process large number of similar samples in parallel. These developments are not directly applicable to the biological defence objective, for which smaller numbers of samples need to be screened for a range of agents. However, some of the method improvements may be of value in a biodefence or arms control setting. For example, the closed tube assay format reduces the workload by removing many of the tedious steps, but at the same time it reduces the risks of sample contamination and thus improves the forensic utility of the technique.

45. Advances have been made in the detection and identification of closely related organisms to strain level in order to distinguish between, for example, virulent and non-virulent strains. Several gene probe methods have been reported for strain typing. Methods fall into two groups: those based on random approaches to identify differences and which require no prior knowledge of the organisms, for example random amplified polymorphic DNA and repetitive extragenic palindromic sequences analyses; and methods that target specific hypervariable regions, for example variable number tandem repeat analysis and insertion sequence analysis.

46. Detection methods are being developed which detect genetic targets indicating the presence of a natural or engineered pathogen. Genetic targets being studied include sequences related to common infection mechanisms and pathways, and sequences common to virulent pathogens (for example pathogenicity islands, antibiotic resistance). Another type of target could be genes that confer environmental stability to organisms, such as sporulation genes, and genes for capsules and resistance to desiccation. The presence of genetically engineered

organisms has also been detected by identifying sequences specific to commonly used cloning vectors.

47. Although antibody technology has progressed steadily over the preceding five years with antibody based biosensors becoming significantly smaller, such technology is still not truly man-portable. The UK national biological defence research and development programme has developed the Prototype Biological Detection System (PBDS), which utilises two antibody-based biosensors for environmental aerosol monitoring. Antibody based detection of toxins and confirmation for a panel of candidate BW agents has been successfully fielded by relatively inexperienced operators.

48. The majority of antibody based detection techniques utilise monoclonal murine immunoglobulin G (IgG) or polyclonal IgG fractions from larger animals. Advances in *in vitro* production of monoclonal murine IgG have been made over the last 5 years, partly because of increasing ethical and regulatory pressures in some countries for a ban on production in ascites tumours. Although hollow fibre technology for IgG production is not a new phenomenon, work has progressed to improve and further understand the secretion of IgG from hybridoma cell lines within these bioreactors. Bioreactors based on cell and product compartmentalisation via molecular weight cut off membranes have been developed and offered as alternatives to ascites production for the sub- 500 mg production market. These systems have reduced the cost of laboratory scale IgG production.

49. Expectations of a tightening of governmental regulations for the export of products containing bovine material have stimulated improvements in media for serum free antibody production. Many assays based on IgG previously produced by ascites may need to be re-optimised as users report a change in characteristics of IgG produced *in vitro*, and perhaps again later when there is a move to serum-free production. Attention is also being given to the use of recombinant technology to move away from production in animal cell lines, and there has been reported expression and purification of fully bioactive whole IgG in plant cells. A major step forward will be the expression of bioactive whole IgG from yeast or bacterial systems as this will make antibody production and purification cheaper, simpler and more reproducible. Single chain antibody can now easily be produced in bacteria.

50. So-called, "Dipstick" hand-held assays based on antigen-antibody specificity have become more widely available. These assays tend to be rugged, readily man-portable devices that have been used to good effect in difficult diagnostic environments. One of the most sensitive designs relies on antibodies adsorbed onto the surface of colloidal gold particles, which give a visual indication of result. These assays have been used with blood samples for the diagnosis of leprosy, bubonic and pneumonic plague, and malarial parasites. Detection of *Bacillus anthracis*

bacteria in environmental samples within 15 minutes using immunogold dipsticks is now possible.

51. Considerable progress in ATP bioluminescence technology has been made over the last five years. In industrial contamination monitoring the trend has been towards simple to use and portable “one-shot” devices containing pre-packaged reagents. UK armed forces now have a generic detection capability based on this principle, providing on-line monitoring of micro-organisms recovered from aerosol particles. The key bioluminescence reagent, the luciferase enzyme, had previously been obtained only from fireflies caught in the wild, but has largely been replaced by a recombinant luciferase that is cheaper and more robust. Adenylate kinase (AK) assays are also being developed. These have allowed a hundred-fold improvement in sensitivity for generic detection, and can be designed as specific assays with limits of detection comparable to PCR. A technique using phage lysis coupled with AK has been developed to specifically detect live bacteria in samples.

52. An optical biosensor, the Resonant Mirror, has been fielded with UK forces. This allows on-line detection of aerosolised toxins. Response times of seconds are possible. A particular appeal is that the logistic burden for this system is low with a single array of detection antibodies immobilised on a sensor surface, allowing 24 hours of continuous monitoring. A matchbox sized surface plasmon resonance (SPR) biosensor is available commercially. Recent developments with a light-scattering SPR biosensor have shown the feasibility of multiplexed real-time detection of bacteria and viruses as well as toxins.

53. The last five years have thus seen the application of a wide range of principles which increase the accuracy, speed, portability and ease of use of detection techniques for field use in detecting biological attack, or for the subsequent identification of agents and toxins in environmental samples. Many of these developments also increase the potential for satisfying the high standards of forensic probity needed for any future sampling and analysis to increase transparency and even to address possible non-compliance concerns under the BTWC.

#### Human infectious disease patterns

54. Despite the energetic international response to the HIV/AIDS pandemic, HIV continues to spread. According to the WHO, AIDS is now the leading cause of death in Africa, and the fourth worldwide. Increasing incidence of AIDS in developing countries where tuberculosis (TB) is epidemic is expected to lead to significant new public health burdens.

55. There continue to be spasmodic outbreaks of emerging and re-emerging infectious diseases with a high morbidity and/or mortality. There have been outbreaks of Ebola in Uganda and neighbouring countries, and cases of Hantavirus in the Balkans. A new viral disease affecting

horses, but communicable to man, was described in Australia and named Hendra. There have been very few human cases to date. Another emerging virus, named Nipah virus after the town in which it was first identified, killed over 100 people in outbreaks affecting pig farmers in Malaysia in 1999. The human cases were initially diagnosed as suffering from Japanese encephalitis, which causes similar haemorrhagic effects. The virus is thought to jump from pigs to humans, and hundreds of thousands of pigs were slaughtered to halt the spread of the virus. There was no evidence of human-to-human transmission, but the US Centers for Disease Control and Prevention classified the virus as a P4-pathogen.

56. There have been some significant outbreaks in developed countries. In New York there was an outbreak of disease caused by West Nile virus, apparently imported in an exotic bird. A rise in the incidence of tularemia in the Balkans has been put down to changes in the population of small mammals that act as the reservoir. Verocytotoxin producing *Escherichia coli* - the toxin is similar to shigatoxin - has a growing prevalence in animals and the environment, a development of concern given the selective impact of this pathogen on children and the old. The virulent *E. coli* strain 0157:H7 has been implicated in a number of outbreaks associated with a high mortality in humans. In spite of a number of surveys, it is still not clear how widespread it is in the environment. Sensitive tests that detect the *E. coli* in cattle have led to calls for more stringent cleansing methods for carcasses in abattoirs. Another food poisoning bacterium that continues to be difficult to control is *Campylobacter*, and in the UK it is now the single biggest identified cause of foodborne illness. Changes in the social structure in some countries, with the majority of armed forces now coming from urban areas, may result in populations with less natural immunity to diseases such as Q fever and psittacosis.

57. It is thus apparent that, even with well-developed public health regimes that can act quickly to detect and control outbreaks, human populations remain susceptible to natural disease outbreaks from a range of diseases. It follows that the chances of human populations being seriously affected by attack with a number of candidate BW agents remain high.

58. There have now been over 100 cases of variant Creutzfeldt-Jacob Disease (CJD) in the UK, and thus it has not (yet) reached the epidemic proportions that some scientists had feared. Increased knowledge about the natural production of prions, and the new molecular biology techniques for modification and enhanced production of proteins, have undoubtedly increased its potential for consideration as a warfare agent. The UK considers it important to recognise that the misuse of prions would contravene Article I of the Convention.

### Smallpox destruction

59. Following the successful completion of the WHO's global project for the eradication of smallpox in 1980, debate continued about the need to maintain stocks of smallpox even if in a limited number of maximum containment (P-4) laboratories. Such stocks were argued to pose a potential attraction for a terrorist group, who might wish to use the virus against unprotected populations. By 1994, the WHO Ad Hoc Committee on Orthopoxviruses had reached consensus that the remaining stocks of virus, at the Centers for Disease Control and Prevention laboratories in Atlanta and the Russian State Research Centre of Virology and Biotechnology (VECTOR) in Koltsovo, should be destroyed. Concerns were however subsequently raised *inter alia* about the lack of an anti-viral agent for use in the event of a smallpox outbreak involving illegal stocks or overlooked natural sources, and that the current smallpox vaccine may not be safe for immunodeficient individuals. The 1999 World Health Assembly reaffirmed the decision to destroy the remaining stocks of virus, but authorised retention until 2002. It was also agreed that a further research priority programme be drawn up and carried out under the strict control of the WHO, and this research is now underway. Some observers have expressed concerns that smallpox cultures may be held outside the two designated laboratories.

### Drug resistance.

60. Methicillin resistant *Staphylococcus aureus* (MRSA) has become a bacterium of great clinical significance in the US, Europe, and Japan, causing infections through open wounds that do not readily respond to antibiotic based treatment. The key genetic component of the resistance is not native to this bacterium, but the heterologous source of the resistance gene is not known. Studies of the genetic profiles and serotypes of strains recovered from outbreaks in several countries strongly suggest that the lineage extends back to the MRSA strains first reported in the early 1960s. MRSA is known to be a great problem in hospitals, but little is known about the incidence of antibiotic resistant bacteria in the normal population. Fears are now spreading that some new-born babies may have suffered from toxic shock syndrome caused by the bacteria's toxin. Recent UK studies reported multi-drug resistant bacteria in apparently healthy children and also that there was no reduction in levels of resistance to sulphonamide antibiotics despite a 45 fold reduction in the use of these antibiotics between 1991 and 1999.

61. Tuberculosis is a disease where multiple drug resistance is a recognised problem. The effects of the combination of AIDS and TB in some population groups is expected to be exacerbated by the multi-drug resistant TB strains which are evolving in prisons and poor communities when antibiotic regimes to treat standard TB cannot be carried through for the several months necessary. (There are several projects to develop a vaccine more effective than the 80 year old BCG vaccine - which often leaves more than 20% of a population unprotected).



Drug resistance and the increasing world incidence of infectious diseases such as AIDS, malaria and TB has been cited to argue that tackling infectious disease should remain a priority and that funding should not be diverted to other pressing health needs, for example to address the rising incidence of chronic non-communicable disease in urban areas.

62. A technique that became outdated once antibiotics were developed but which may receive renewed attention in the future as a means to deal with antibiotic resistant bacteria is the use of bacteriophage (virus that infects a bacterium). One current disadvantage of known bacteriophages is that their use would depend on prior diagnosis of the bacterial strain.

63. The existence of so many naturally multiple drug resistant strains could greatly complicate the therapy of secondary (natural) infections in BW casualties.

#### Disease in agriculture

64. The globalisation of trade has brought with it a growing sensitivity of importers and consumers to the risks of animal and food borne diseases. The huge trade impact of outbreaks of disease in agricultural animals, with foot and mouth disease (FMD) of cattle as the foremost example, is clear to all. One major problem is the export of meat and other agricultural products for personal consumption or local sale, not declared to the authorities of the importing country. Such products can carry exotic human, animal or plant disease that can have a devastating effect. Though the cause of the 2001 outbreak of FMD in the UK may be difficult to establish with certainty, it is likely that the virus came into the UK through illegal importation of meat. The increased movement of people and agricultural products has also exacerbated the spread of economically damaging plant disease. Increased exportation of exotic plants, often not declared, has considerably complicated the job of plant regulatory authorities. In both animal and plant disease, measures which prohibit trade with countries when a particular disease is reported leads to the concept of pest free areas, which demands survey work which can only be performed if there are internationally standardised analysis techniques used in accredited laboratories.

65. According to OIE statistics, 66 countries have reported infectious disease outbreaks due to emerging animal diseases during the last five years. Increasing attention is being given to mitigating the adverse effects of emerging animal diseases on livestock production, trade and public health. OIE initiatives include additional reference laboratories, revised international health codes (these are the standard texts used to promote the harmonisation of regulations for trade in animals and animal products), and co-ordinated international training programmes for managers of veterinary services.

66. Difficult weather conditions and persistent civil strife have caused exceptional food shortages in sub-Saharan Africa. On top of this, there is a growing realisation that many plant

pests, which cause problems for trade within tropical or sub-tropical zones also, can threaten European countries through trade (and tourism) links. In the reverse direction, for countries that have diversified by fruit, vegetables and ornamentals production under the cooler conditions at high altitudes, there is a growing risk of damage from pests introduced from temperate regions.

67. The economic impact of Newcastle Disease and its effect on trade in commercial poultry in developed countries is well known, but experts also recognise the problem of the endemic presence of the disease in developing countries and the effect on village chicken production. Efforts have been made to introduce vaccination programmes in the form of heat resistant live vaccines, but experts advise that vaccination cannot resolve the problem unless matched by education programmes into disease spread and poultry husbandry.

68. Airborne spread is not generally regarded as being implicated in outbreaks of Newcastle Disease reported in recent years, and there has nearly always been a alternative, more likely, cause, particularly the movement of poultry and humans. For instance, in 1997 there were 38 outbreaks confirmed in poultry in the UK between January and April. Nucleotide sequencing and phylogenetic analysis showed close similarity between the UK isolates and virus responsible for outbreaks in Scandinavian countries the previous year, including a isolate from a feral goosander. Unusual patterns of movement of migratory birds at the end of 1996 and beginning of 1997 suggest that this may have been the vehicle for introduction of the virus into Britain. New knowledge of the molecular basis of pathogenicity of NDV have led to an OIE definition which differentiates virulent from avirulent virus on the basis of differences in a region of the F (fusion) protein of the virus. PCR tests based on this genetic difference however still need improvement.

69. Major outbreaks of avian flu in poultry with spread to numerous sites and huge economic losses continue to be reported, and occurred in Hong Kong in 1997 and Italy in 1999/2000. The high cost of vaccination, since only inactivated vaccines are available, means there is economic pressure to stop vaccination once the immediate threat is thought to have passed. Of the 18 outbreaks of Highly Pathogenic Avian Influenza between 1959-2000, in only two was vaccination used, and then in combination with increased biosecurity measures. The other 14 outbreaks were controlled by a rigorous slaughter and biosecurity regime. Although biosecurity measures in poultry farming are often regarded as costly and laborious by those involved, experts recommend that they are an essential investment as a means to prevent the risk of introduction of avian flu and ND, as well as other endemic viruses that may affect the birds and reduce their yield.

70. An outbreak of *Thrips palmi* at a commercial site in the UK has been investigated, and an eradication programme completed.

71. In spite of some improvements in trade regulations for animal and plant products, and in diagnostic capabilities, the magnitude of the effects of natural disease outbreaks especially for animal diseases and the public awareness of this may well have increased the attractiveness of anti-animal and anti-crop BW to potential proliferators.

#### Pest control in agriculture

72. The peaceful use of micro-organisms as agricultural pesticides has expanded in developing countries since the last Review Conference, and the use of transgenic plants that have increased pest resistance is likely to be the most important innovation since the appearance of chemical pesticides in the 1940s. The genetic modification approach is seen by many as the best method for delivering the benefits of new crop technologies to farmers especially those in developing countries, where agricultural supplier and advice networks are often inadequate to channel new information, chemicals or equipment.

73. Genetically modified crops are being grown on a large scale in commercial agriculture in several countries, notably the United States, China, Canada and Argentina. In one type of modification, genes for an insecticide toxin of *Bacillus thuringiensis* (Bt) are inserted into crops. This modification allows the plant to produce the toxin and thus kill insects that feed on it. Data for 2000 indicates that, of the 29 million hectares of genetically modified crops grown in the USA, Argentina and Canada, 24% was transgenic maize (corn) or cotton hybrids containing *cry* genes derived from Bt. China, Mexico and Australia are also major growers of these Bt crops, and commercial production in India is planned. In many cases Bt crops allow use of chemical pesticide to be reduced, and there have been reports of significant increases in yields and profits. The chemical pesticides are still needed to control populations of secondary pests, such as aphids, plant bugs and thrips.

74. Direct application of Bt formulations as an insecticide has continued, but its share of the world pesticide market has been stable at around 1-2% in the last decade. Bt is a naturally occurring soil bacterium, and although not regarded as a human pathogen there is a report of the isolation of the bacterium, though not a serotype used in agriculture, from the wound of a severely injured soldier. Potential concerns about the risks to human health arising from the conventional spraying of Bt formulations, a technique in use as long as 40 years ago, have stimulated a very large number of laboratory and epidemiological studies, and more recently such studies have addressed the possibility of allergic responses to Bt transgenic plants. Increases in human antibody levels following exposure to Bt products have been reported, but there was no increased incidence in asthma or other illnesses, and many experts conclude that there is no reason to doubt the safety of Bt insecticides.

75. There has been increasing attention to the development of other pesticides produced using fermentation techniques, though no new products are yet in routine use. The potential for use of fungi in the field as opposed to glasshouse situations has increased as a result of improved formulation technology. The constraint of high humidity has been overcome in some cases, facilitating the use of fungi as insecticides in dry climates such as deserts, for example for locust control. An advance with major significance for crop protection is the demonstration of the use of fungi to produce the spinosyn group of insecticides. These molecules have high activity against pests that are resistant to conventional chemical insecticides, but appear to be very safe for man. The avermectin insecticides that are produced by *Streptomyces* have been used for some time but appear to have more limited potential.

76. A continuing problem that is deemed to erode confidence in microbial based control agents is the variation in performance in the field caused by variable quality of the formulations used. The lack of quality control in manufacture, even when low technology methods are used, damages the reputation of the technology and also may pose health risks to workers in production plants or to people exposed in the field. Experts in developed countries are contributing to initiatives to improve manufacturing quality control worldwide.

77. In the past five years there has been progress in developing commercial bioinsecticides in the US, EU, India and South Asia which make use of insect viruses primarily of the nucleopolyhedrosis viruses (NPV). Overseas aid from the UK has funded projects to develop an NPV industry in India, and viral pesticides to control army worm and diamond back moth in parts of Africa. Currently, NPV is produced commercially *in vivo* in several countries worldwide, and the future will see *in vitro* production in fermenters of up to 20,000 litres.

78. Some of the above technologies, including microbial manufacturing plants and field delivery systems, could be misused in illicit BW programmes.

Global initiatives to tackle disease.

79. The threat caused by the increasing globalisation of disease has been recognised by the key international bodies concerned with addressing human, animal and plant disease. By the turn of the millennium, a number of important global initiatives and partnerships in disease prevention and control had been created. These are relevant to Article X of the Convention.

80. The 1995 World Health Assembly decided to revise and update the WHO International Health Regulations, the legal framework for WHO's alert and response activities. As the first step in the revision process, the idea to replace disease-specific reporting with a system of syndromic reporting was field tested. The scheme was not taken forward because of difficulties

in reporting syndromes in the field, and because syndromes could not be linked to preset rules for control of spread. There are still only three diseases for which reporting is mandatory to the WHO, cholera, plague and yellow fever.

81. WHO figures indicate that HIV has now infected 58 million men, women and children, with 22 million deaths. It is recognised that a safe, effective vaccine suitable for use especially in developing countries is badly needed to complement existing prevention strategies. WHO and the UN Programme on HIV/AIDS (UNAIDS) have joined forces to establish a new HIV Vaccine Initiative.

82. The WHO Vaccine Preventable Diseases Monitoring System collects, compiles and disseminates data officially reported by WHO member states, on the incidence of vaccine preventable diseases and on immunisation coverage. There are a number of significant international initiatives to accelerate the introduction of new and under-used vaccines, aiming to address the practical difficulties of strategy development, logistics, funding, supply and quality control. The UN has set up a Global Health Fund to tackle HIV, malaria and TB, and the manufacturers of the expensive TB drugs have agreed with the WHO to supply at cost price to poorer countries. The Global Alliance for Vaccines and Immunisation is an alliance which includes governments, intergovernmental bodies such as UNICEF, WHO and the World Bank, philanthropic institutions, and trade associations. With resources for 2001 -2005 in excess of US \$ 1 billion, the Fund will help to provide new and under used vaccines and the means for delivery in the field, as well as funding to help governments strengthen their basic immunisation services. Under utilised vaccines include those for rubella and yellow fever, and the new vaccines being considered include hepatitis B, *Haemophilus influenzae* type b (Hib), and the pneumococcal conjugate vaccines.

83. After the successful eradication of smallpox, efforts continue to eradicate polio by 2005, concentrating on the 20 or so countries where it still occurs. Objectives include obtaining sufficient finance and political support, and particularly to secure access to immunise all children including those in areas affected by conflict.

84. The WHO has recognised that trade is often adversely affected when certain public health risks occur, and has set up links with the WTO Committee on Sanitary and Phytosanitary Measures. Other new joint initiatives have included WHO, FAO and OIE consultations on BSE (bovine spongiform encephalopathy), that considered issues of public health, animal health and trade. The BSE talks recommended that BSE and the human disease variant CJD should be considered as an international issue because potentially infected BSE materials have been distributed throughout the world through trade of live cattle, cattle products and by-products. Countries should take care not to be complacent about their risk from BSE. The meeting stated

that the extremely low initial incidence and limited clustering of BSE cases, protracted latency and non-specific nature of the early clinical signs of BSE tend to mask the severity of the problem.

85. An OIE/FAO joint conference in April 2001 on FMD passed resolutions which *inter alia* recognised that strengthening of veterinary services and the creation of international control and eradication programmes for epizootic (animal) diseases was not only in the interests of developing countries but also would reduce the risk of spread of such diseases worldwide.

86. The Global Rinderpest Eradication Programme has a target to eradicate rinderpest worldwide by the year 2010. Only three small reservoirs of rinderpest remain, in Asia and Africa, but FAO experts are concerned that there is now a high vulnerability to epidemic resurgence, which demands renewed attention to elimination of the last reservoirs of disease and management of the cessation of vaccination.

87. Since the 1995 Agreement on the Application of Sanitary and Phytosanitary Measures of the WTO came into effect, many southern countries have found difficulties in meeting the extra burden and resource implication for their phytosanitary services. The 1997 revision of the International Plant Protection Convention (IPPC) emphasises the International Standards for Phytosanitary Measures and the procedures of Pest Risk Analysis (PRA), which allow import decisions to be based on scientific evidence. However, the revised IPPC has not yet received sufficient signatures to come into effect, and the lack of meaningful international guidelines for PRA five years on is causing concern. Regional Plant Protection Organisations have however been set up as co-ordinating bodies in the various geographic regions, and to gather and disseminate information and promote standards.

Molecular biology applications and crops.

88. New arthropod targeted transgenic crops and trees are being developed: this includes a transition from single to multiple Bt toxin genes, and use of entomopathogens such as bacterial *Xenorhabdus* species and the *Heliothis* stunt virus. Bt toxin expression can be increased markedly by engineering the toxin genes into the chloroplast, where toxin yields can amount to up to 50% of the chloroplast proteins. Despite the enthusiastic uptake of such technology, there are concerns regarding the potential adverse effects to human health and to the environment, and its application in some countries has been subject to government scrutiny and regulation.

89. There have been a number of projects to increase the efficacy of viruses as insecticides by incorporating genes for toxins, diuretic hormones or other genes that enhance the virus killing power. The engineering of scorpion toxin into baculoviruses is well known approach, and other

polypeptides are also being considered - for example, peptides from the venom of the parasitic wasp *Bracon hebetor*. Recently, genetic modification has been used to increase the pathogenicity of baculovirus.

90. Genetic modification technologies are likely to become the primary route to producing new varieties of plants with resistance to diseases. An increasing number of projects involve incorporation of viral genes, using the promoter sequences of the virus in conjunction with the coat or movement proteins genes to build in virus resistance. Environmental impact needs careful consideration because such genetic engineering results in incorporation of viral genomes throughout the tissues of the plant, which could create opportunities for recombination with wild viruses leading to unwanted changes in their pathogenicity, host range etc.

91. Genetically modified potatoes with resistance to potato virus Y and potato leaf roll virus have been developed, but are currently not believed to have significant commercial appeal because of public reticence about engineered foods. The UK government and other international donors are funding the development of rice with resistance to yellow mottle virus by incorporating RNA polymerase genes into the rice.

92. Gene silencing is a technique that makes use of the plant's natural anti-viral defence system. A systemic signal molecule is produced in the plant, along with short RNA molecules that are homologous to the target in the host DNA. This methodology has been developed at the Sainsbury Laboratory in the UK and is marketed. Plant virus vectors are used; they carry fragments of host plant DNA throughout the plant as they replicate, and the gene in question is silenced.

93. There have been great improvements in antibody based diagnostic tests for plant viruses, and dipstick tests developed in the UK for a number of viruses are now used in the field by plant health inspectors. Antibody tests are also used to monitor Bt toxin expression in genetically modified crops.

94. Genetic typing has also improved the direction of classical techniques for breeding disease resistant crops. An example is a attempts to reduce the impact of more virulent strains of the potato fungus *Phytophthora infestans* which have reduced yields world-wide in recent years. New varieties of potatoes have been bred by crossing with Peruvian wild type potato subspecies *Solanum tuberosum andigena* to make use of its resistance genes.

95. Genetic typing (genetic fingerprinting) carried out in investigations of animal and plant disease outbreaks provided new insights into the relationships between outbreaks in different

countries and over a period of time. An example for the typing of Newcastle Disease in birds has been given above. An example for plant disease comes from the recent published finding that the strain of the fungus *Phytophthora infestans* which caused the Irish potato famine in the 19th century lacks a gene sequence found in the modern strain 1b, and this overturns the classical theory that 1b is directly descended from the Irish strain.

96. New insights into the interactions between micro-organisms and plants unfortunately will also increase the potential for design of anti-crop agents with improved properties such as a broader infection window in the life cycle of the plant. On the other hand, advances in diagnostic capabilities should make it easier to distinguish natural and unnatural outbreaks of infectious disease.

#### Trends in protein production technologies

97. The huge increase in the number of biotechnology ideas in the research stage has not been matched by an increase in the manufacturing capacity for the ultimate products, and in developed countries this has led to significant queuing times for the limited contract manufacturing slots. Very few new products are made by extraction from natural sources. A factor contributing to this bottleneck is the change in type of protein product in development, to antibodies - made by growing tissue culture cell lines, often in fermenter vessels. In the 1980s many drugs based on monoclonal antibodies failed to reach their potential because their repeated use led to immune reactions to the foreign protein. A number of antibody based drugs recently launched may have largely circumvented this problem by using humanised antibody from genetically transformed cell lines or murine/human monoclonal chimeric antibody. A current estimate is that there are more than 70 monoclonal antibodies in various stages of clinical development, amounting to about 20% of all biotech drugs in development. These large and complex proteins are often poorly expressed, difficult to recover and purify, and the typical dose is up to a hundred-fold more than for a hormone or enzyme. Their production demands large fermentation and downstream processing systems. Of the licensed biological products that are produced in cells, about half come from micro-organisms and half are made in cell lines of human or animal origin.

98. The increased experience in production of antibodies suitable for human administration has increased the potential for manufacture of therapeutic antisera in dealing with significant outbreaks of bacterial disease for example after a BW attack.

99. Early ideas for pharmaceutical production in transgenic animals or plants have been slow to come to fruition. Transgenic animals may yet have considerable potential for production of high volume products, for example in cows' milk. Disadvantages are long development times, and difficulty in ensuring and demonstrating the absence of risk from transmissible spongiform encephalopathies caused by prions. Transgenic plants could also provide large quantities of



material, but aberrant glycosylation could complicate the development of some products. Tobacco mosaic virus is often used as a virus vector: it spreads rapidly and systemically in the plant to give high yields of protein, and the fact that it is not seed or pollen transmitted and not vectored by insects gives a low risk of unwanted spread from the host crop. Using this technique, companies in the US claim to have produced experimental batches of hundreds of different proteins from ton quantities of infected crops, and there are reports of plans to scale up to thousands of acres. Avian transgenics may also be promising as a production vehicle: the production of eggs on a large scale is already well organised world-wide in vaccine industries; sterile separation methods are well developed; and glycosylation is unlikely to be a problem. Any of these new production routes could however, also be misused to produce significant quantities of proteins or peptides for use in BW.

#### International co-operation and biosafety: activities under the Biodiversity Convention

100. The Convention on Biological Diversity, one of the main outcomes of the 1992 Rio Earth Summit, is now well established and increasing its impact. The UK is the fifth largest donor to the Global Environmental Facility, a financial mechanism set up under the Convention to provide resources and expertise for projects which assist developing countries in biodiversity conservation and sustainable use. The UK also supports such work bilaterally and through the UK's Darwin Initiative. An example of a microbiological project recently completed under the latter is the three year Darwin Project on Caribbean Fungi. This was intended to raise the profile of fungi in the Caribbean, by using Cuban expertise and developing Trinidad and Tobago expertise. Scientists from the UK's International Mycological Institute delivered equipment, conducted field work, and trained staff in Cuba and Trinidad. National strategies were developed, a guide to the identification of microfungi on sugar cane was produced, more than 9 data bases developed, and hundreds of culture collections were enhanced. The infrastructure for a regional identification service is now in place.

101. The Cartagena Protocol on Biosafety to the Convention was agreed in January 2000. It aims to guard against risks to biodiversity arising from transboundary movement of living modified organisms (essentially, genetically modified organisms). Provisions under the Protocol introduce requirements prior to export of "living modified organisms" (LMOs) including commodities for food, feed and processing. An intergovernmental committee is currently developing arrangements for implementation. Key aspects of the Protocol are public access to science and information and public participation in debate on biosafety issues. A publicly accessible, Internet-based "Biosafety Clearing-House" will enable the exchange of scientific and other information about traded LMOs, while a "roster of experts" will give Parties access to scientific expertise to make informed decisions on the import of LMOs. The Protocol also requires Parties to consider, in consultation with other relevant international bodies, the potential elaboration of standards on the identification, handling, packaging and transport of LMOs.

Parties are also due to consider the possible need for rules and procedures on liability and redress for damage resulting from transboundary movements of LMOs. Finally, the Protocol places an emphasis on “capacity building”, whereby developing countries are assisted in developing the necessary legal, institutional and technical frameworks for effective implementation of the Protocol.

#### Means of delivery of agents or toxins

102. While there has been work on developing new application techniques for microbial pesticides including electrostatic sprayers, their uptake has been very limited. Most application of microbial pesticides is still through traditional hydraulic or spinning disc systems. However, the potential for effective delivery of biological agents or toxins as weapons has undoubtedly increased because of the increasingly widespread availability of long range missiles, sub-munitions and high specification multi-barrel rocket launchers which could be modified to deliver biological weapons.

#### Use of pathogens to control weeds and ‘criminal’ crops

103. Over 100 microbial pathogens of plants have been identified as potential biocontrol agents for weeds, but only a handful have entered commercial use and then only on a limited scale. Examples are the biocontrol of tough grasses, and a fungus that prevents weed growth from tree stumps. Practical difficulties typically include slow speed of action and poor selectivity for host.

104. A number of countries are involved in supporting the UN Drug Control Programme in projects to develop pathogens for release as a specific bioherbicide against ‘criminal’ drug crops such as coca, opium and marijuana. The fungus *Fusarium* is being considered for use against coca plants from which cocaine is manufactured, and another fungus, *Pleospora papaveraceae*, is being considered to attack opium poppies. NGOs have raised concerns about whether the possible effects of persistence in the environment including unforeseen genetic transfers or infections of other plants are being thoroughly addressed. The UK believes that the use of biological agents or toxins for control of criminal crops should be discussed at the forthcoming Review Conference.

#### Bioremediation: the destruction of materiel

105. Studies of the potential for the release of specific micro-organisms into the environment to destroy pollutants continue. An example is the destruction of residues of ethylene diglycol (antifreeze) after its use on airport runways and aircraft. Critics of this approach have been concerned about the potential for gene transfer to the gene pool of natural organisms. Field trials attempting to address such concerns have included the release of bacteria into which had been

engineered the marine bacteria bioluminescence *lux* gene, allowing the spread of the trial bacteria to be monitored with a hand held light detector. While no microbial bioremediation methods have been adopted for the large scale destruction of chemical agent in national munition destruction programmes related to the provisions of CWC, interest has continued in the potential of micro-organisms for neutralising residues of CW agents in environmental samples. For example, micro-organisms capable of utilising thioglycol (a breakdown product of mustard) have been isolated.

106. Bioremediation technologies clearly have the potential for development as a means of warfare or for hostile use against materiel crucial for normal civilian life or military operations, such as oils, rubbers and plastics.

#### Countering the threat of BW terrorism

107. In the wake of the Aum Shinrikyo's attempts at bioterrorism in 1995, several countries have taken significant measures to increase their ability to counter the use of biological agents by terrorists. Measures include increased funding to study diseases affecting human or animals and to develop pathogen and toxin detection devices; planning for procurement and stockpiling of vaccines to protect armed forces and civil populations; and regional and national contingency planning and training which co-ordinates the activities of civil emergency and police services, health authorities and specialist scientist advisers. In the United States, the objective of preventing unauthorised attempts to obtain pathogens and toxins from legitimate laboratories was one of the rationales for a new national system to register laboratories and licence the acquisition and transfer of 'select agents' on a list, with appropriate research and clinical exemptions. Also in the US, the Advisory Committee on Immunisation Practices in February 2001 produced new draft guidelines on the use of smallpox vaccine in the event of an outbreak of smallpox, in the context of bioterrorist use of the agent. Government studies to assess the threat of bioterrorist attacks have included a Canadian study of the threat of use of FMD virus, and a Japanese study which listed 20 bacterial and virus species, including smallpox.

108. National anti-terrorist programmes where significant funding is being applied to the procurement of detection and medical countermeasures may also be expected to improve national preparedness against BW attacks by other states.

#### Impact of the entry into force of the CWC

109. There has been a major increase in industrial uses of micro-organisms or their extracted enzymes to catalyse single steps in chemical manufacturing processes otherwise based on conventional chemistry. These applications, known as 'biotransformations', mostly focus on production of intermediates where a directed chirality is required. This reflects a large increase

in the number of small molecule drugs where chirality is realised to be important for the therapeutic effect. When whole micro-organisms are used they are often produced on the chemical production site, but isolated enzymes are often bought in from specialist manufacturers. Facilities using biotransformation are likely to be encountered in future plant site declarations under the Convention on the Prohibition of the Development, Production, Stockpiling and Use of Chemical Weapons and on their Destruction (CWC).

110. The CWC, which entered into force on 29 April 1997, in effect precludes the development, production or use of chemicals as weapons against man or animals. The so-called general purpose criterion of the CWC covers any toxic chemical or its precursor, regardless of its origin or method of production, and so there is considerable overlap with the BTWC. Two toxins, ricin and saxitoxin, are included in the Schedules of chemicals, and as they are in Schedule I of the Verification Annex they are subject to licensing and declaration provisions, and limitations on transfers, which are permitted only between States Parties. The Schedule I entries for these two toxins are limited by CAS (Chemical Abstracts Service) numbers, which for saxitoxin does not cover a number of forms of the toxin that could be of concern. To the UK's knowledge, no uncertainties have been identified about either of these toxins during inspections of Schedule I facilities by officials of the Technical Secretariat of the Organisation for the Prohibition of Chemical Weapons. The CWC also provides for declaration and inspection of certain chemical production facilities not producing Scheduled chemicals, under the category of Other Chemical Production Facilities (DOC facilities). The UK is not aware that any facilities producing chemicals by use of fermentation technology, whether toxins or any other molecules, have been declared as DOC facilities, nor does this seem likely at present given the DOC declaration threshold of 200 tonnes (30 tonnes where the chemical contains the elements phosphorus, sulphur or fluorine). In our view, any measures, which may evolve under future agreements to strengthen the BTWC, and which are designed to provide transparency for national activities, should involve many more toxins than the two covered by the Verification Annex of the CWC.

---